



It's time to listen: there is much to be learned from the sounds of tropical ecosystems

Jessica L. Deichmann^{1,16} , Orlando Acevedo-Charry^{2,3}, Leah Barclay⁴, Zuzana Burivalova⁵, Marconi Campos-Cerqueira², Fernando d'Horta⁶, Edward T. Game⁷, Benjamin L. Gottesman⁸, Patrick J. Hart⁹, Ammie K. Kalan¹⁰, Simon Linke¹¹, Leandro Do Nascimento¹², Bryan Pijanowski⁸, Erica Staatterman^{13,14}, and T. Mitchell Aide^{2,15}

¹ Center for Conservation and Sustainability, Smithsonian Conservation Biology Institute, National Zoological Park, Washington, DC, USA

² Sieve Analytics, San Juan, PR, USA

³ Colección de Sonidos Ambientales, Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, Colombia

⁴ Queensland Conservatorium Research Centre, Griffith University, Nathan, Qld, Australia

⁵ Woodrow Wilson School of Public and International Affairs, Princeton University, Princeton, NJ, USA

⁶ Graduate Program in Genetics, Conservation and Evolutionary Biology, INPA, Manaus, AM, Brazil

⁷ Global Science, The Nature Conservancy, Brisbane, Qld, Australia

⁸ Department of Forestry and Natural Resources, Purdue University, West Lafayette, IN, USA

⁹ Department of Biology, University of Hawaii at Hilo, Hilo, HI, USA

¹⁰ Department of Primatology, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

¹¹ Australian Rivers Institute, Griffith University, Nathan, Qld, Australia

¹² Department of Wildland Resources and Ecology Center, Utah State University, Logan, UT, USA

¹³ Bureau of Ocean Energy Management, Office of Environmental Programs, Sterling, VA, USA

¹⁴ Beneath the Waves, Inc., Herndon, VA, USA

¹⁵ Department of Biology, University of Puerto Rico, San Juan, PR, USA

ABSTRACT

Knowledge that can be gained from acoustic data collection in tropical ecosystems is low-hanging fruit. There is every reason to record and with every day, there are fewer excuses not to do it. In recent years, the cost of acoustic recorders has decreased substantially (some can be purchased for under US\$50, e.g., Hill *et al.* 2018) and the technology needed to store and analyze acoustic data is continuously improving (e.g., Corrada Bravo *et al.* 2017, Xie *et al.* 2017). Soundscape recordings provide a permanent record of a site at a given time and contain a wealth of invaluable and irreplaceable information. Although challenges remain, failure to collect acoustic data now in tropical ecosystems would represent a failure to future generations of tropical researchers and the citizens that benefit from ecological research. In this commentary, we (1) argue for the need to increase acoustic monitoring in tropical systems; (2) describe the types of research questions and conservation issues that can be addressed with passive acoustic monitoring (PAM) using both short- and long-term data in terrestrial and freshwater habitats; and (3) present an initial plan for establishing a global repository of tropical recordings.

Key words: conservation technology; ecoacoustics; passive acoustic monitoring; soundscape.

“The universe is your orchestra. Let nothing less be the territory of your new studies” Raymond Murray Schafer (1969)

In an era of rapid environmental change, remote sensing methods are particularly important for ecology and conservation biology because they produce consistent data streams that can be analyzed over different spatial and temporal scales (Kerr &

Ostrovsky 2003; Nagendra *et al.* 2013; Turner *et al.* 2003). Passive acoustic monitoring (PAM) is one way to characterize and evaluate ecosystems remotely using sounds. First developed for use in the marine realm (Tavolga 2012), autonomous recordings can detect a range of sounds produced by natural and physical phenomena (Krause 1987). The ‘soundscape’ includes all sounds emanating from any given habitat, which can be classified with respect to their source: geophony (climate and geography), biophony (all wildlife), and anthrophony (human activities;

Received 21 February 2018; revision accepted 8 June 2018.

¹⁶Corresponding author; e-mail: deichmannj@si.edu

Pijanowski *et al.* 2011). Analysis and monitoring of these various contributions to a soundscape can permit rapid assessment of biodiversity as well as the health and stability of an ecosystem (e.g., Blumstein *et al.* 2011, Pijanowski *et al.* 2011, Fuller *et al.* 2015, Bertucci *et al.* 2016, Burivalova *et al.* 2017, Deichmann *et al.* 2017, Staaterman *et al.* 2017).

APPLICATIONS OF ECOACOUSTICS IN THE TROPICS

Many tropical biologists have been startled by the sound of a nearby treefall, while others have been warned of an oncoming storm by croaking toucans or the presence of a predator by screeching squirrel monkeys; yet many of us have never considered that these sounds are data that can be harnessed to answer questions about tropical ecosystems. Here are a few examples of the types of questions that can be answered using sounds:

POPULATION DYNAMICS AND ACTIVITY PATTERNS.—We know very little about natural activity fluctuations within tropical forest communities, and perhaps even less in tropical freshwater systems. Thus, it is difficult to precisely assign causal relationships between human activities and changes in biodiversity (Thompson 2003). For example, is the decline in abundance of a hornbill species in an Indonesian forest a part of a naturally occurring seasonal and superannual fluctuation pattern, or is the population actually decreasing due to hunting, logging, and habitat loss? If measurements are taken during a ‘low’ part of an undetected cycle, small population numbers could make the impact of an otherwise-sustainable hunting practice appear catastrophic. Alternatively, unsustainable hunting rates could be seen as deceptively benign if measurements were taken during a peak time. Recording soundscapes regularly to span the natural cycles of animal activity helps us correctly understand these patterns (Bridges *et al.* 2000, Towsey *et al.* 2014, Linke *et al.* 1999), which otherwise would be extremely difficult to decipher using traditional biodiversity monitoring methods.

BROAD SPATIAL SCALES.—Our current methods for comparing biodiversity of multiple habitats (beta diversity) are insufficient. This task is notoriously difficult in tropical forests and streams due to the sheer number of species present and the amount of sampling necessary. The ability to deploy multiple acoustic sensors across landscapes in a short period of time enables simultaneous recording, which allows researchers to make meaningful comparisons and tackle elusive patterns in tropical forest and freshwater fauna (e.g., Bormpoudakis *et al.* 2013, Gasc *et al.* 2013, Rodriguez *et al.* 2014). For instance, PAM can improve our understanding of ecological processes across entire elevational gradients helping us to track the impact of climate change on animal distributions (Campos-Cerqueira *et al.* 2017).

RAPID INVENTORIES AND SPECIES OF CONSERVATION CONCERN.—The presence of rare and cryptic species in tropical habitats is difficult to detect in short trips to the field (Thompson 2003, Plaisance *et al.* 2011), but PAM methods have been successfully used to

detect such animals in densely forested habitats, producing results that would otherwise require massive search efforts by field crews. For example, PAM has been used to estimate the presence and abundance of African forest elephants (*Loxodonta cyclotis*) inhabiting dense rain forests of Central Africa (Wrege *et al.* 2017) as well as cryptic fish in tropical coastal habitats (Staaterman *et al.* 2017) and an endemic and threatened bird in Puerto Rican mountains (Campos-Cerqueira & Aide 2016). Invasive species such as fish (Rountree & Juanes 2017) and pest insects (Mankin *et al.* 2011) have also been detected using PAM. Likewise, PAM can detect the recovery of species extirpated from a site after natural disaster, disease, or other perturbation (Butler *et al.* 2016). The ability for PAM data to be collected rapidly from many places but analyzed later makes it a valuable tool for rapid inventories (Sueur *et al.* 2008, Ribeiro *et al.* 2017), which tend to be costly and difficult to fund.

HUMAN IMPACTS AND SHIFTING BASELINES.—Comparing soundscapes in areas under different management regimes allows for a rapid understanding of the intensity of impact caused by different human activities (e.g., Alvarez-Berríos *et al.* 2016, Burivalova *et al.* 2017, Deichmann *et al.* 2017). Examples include changes in habitat structure (Tonolla *et al.* 2010, Geay *et al.* 2017) or levels of hunting activity in protected areas (Astaras *et al.* 2017). Furthermore, acoustic data collected over the long-term can be used to answer broader questions regarding the effects of environmental change on species abundance, phenology, distribution (Campos-Cerqueira & Aide 2017, Campos-Cerqueira *et al.* 2017), and behavior (Llusia *et al.* 2013, Narins & Meenderink 2014). For example, acoustic monitoring has been used to demonstrate changes in the seasonal onset of birdsong (Buxton *et al.* 2016), which may be indicative of climatic influences on the timing of reproduction. Acoustic ‘time-capsules’—measurements made in the past or the present—will be critically important for similar observations in the decades to come.

ADVANTAGES OF PASSIVE ACOUSTIC MONITORING

Using PAM, rather than traditional methods, to monitor and analyze biodiversity will help us do a better job of understanding and conserving tropical terrestrial and freshwater ecosystems. Netting, trapping, distance sampling, visual transects, etc. are labor-intensive, expensive, and logistically impractical in many places—often even more so in the tropics than in the temperates. In addition, most observations of animal behavior are influenced by human presence and limited to daylight hours. Crucially, the autonomous nature of acoustic sensors permits continuous collection of PAM data without biases from the ‘observer effect’ (Shonfield & Bayne 2017). PAM can cover broad spatial and temporal scales, including simultaneous and long-term monitoring, which is simply not possible with traditional methods (Linke *et al.* 2018). This can even be done in real time (e.g., Van Parijs *et al.* 2009, Aide *et al.* 2013), providing researchers and managers with information necessary for immediate decision making, and make adaptive management more feasible.

Finally, collection of big data through PAM creates permanent records that can be reanalyzed when new analytical tools become available, when additional research questions arise, or to compare past to present conditions.

The related technique of camera trapping has greatly improved our capacity to estimate species composition, abundance, and density of medium to large-bodied mammals and birds—groups that are difficult to study using traditional methods—in terrestrial (Burton *et al.* 2015) and arboreal habitats (Gregory *et al.* 2014). That said, camera trapping is restricted to this subset of species (but see Hobbs & Brehme 2017) and the detection range is relatively limited. PAM has the additional benefit of having broader detection ranges [e.g., maximum 1 km detection radius calculated for primate sounds (Kalan *et al.* 2015); up to many km depending on frequency, microphone height, and habitat type (Darras *et al.* 2016)] and sampling a wider range of taxonomic groups (Aide *et al.* 2017). We consider camera trapping and acoustic monitoring to be complementary in terms of taxonomic groups and advocate the use of both methods where possible.

CHALLENGES

While PAM holds many advantages over other methods, it would be remiss not to recognize that challenges do exist. For example, as with other methods that result in the collection of big data, PAM faces the challenge of data storage and management. Storing recordings on multiple hard drives is not expensive, but it is not a particularly effective way to encourage their use in analyses by the broader community. Furthermore, extracting meaningful biological information from recordings is complex. Automated detection tools for species-level analyses have advanced significantly over the last decade (e.g., Aide *et al.* 2013, Kalan *et al.* 2015, de Camargo *et al.* 2017). Still, there are limitations to automatic approaches because they can be initially time-consuming and they require training data to create different classifiers for different species as well as programming or signal processing expertise to develop automated species identification models. At the soundscape level, many acoustic indices and soundscape analysis methods have been proposed and used for the assessment of biodiversity (e.g., Sueur *et al.* 2008, Pieretti *et al.* 2011, Villanueva-Rivera *et al.* 2011, Gasc *et al.* 2013, Fuller *et al.* 2015, Vega *et al.* 2016, Aide *et al.* 2017, Rankin & Axel 2017), yet there is no consensus to date as to which are most effective, primarily due to the difficulties in generalizing across taxa and ecosystems (Buxton *et al.* 2018). Existing indices can also be sensitive to geophony such as rain, wind, and river flow, or can be skewed by certain acoustically dominant species (Staaterman *et al.* 2017, Linke *et al.* 1999). Most also lack measures of uncertainty (e.g., detection probabilities)—an issue likely to be exacerbated in highly diverse tropical habitats. Nevertheless, collection of acoustic data now opens up the possibility of analyzing long time series of sounds in the future as analytical methods become more advanced and standardized—a possibility that can only be realized if we start recording now.

BROADER IMPACTS OF ECOACOUSTICS

In addition to serving as permanent records of ecosystem health and providing data for scientific research, animal sounds can serve as an alluring tool for engaging public audiences. Camera trapping has been successful for many reasons, but chief among them is the charismatic nature of the resulting photographs—who doesn't smile when they see wildlife 'selfies' or animals caught in the act of defiling a camera? We argue that sounds can be just as captivating—many of us have seen public audiences become wide-eyed when we play them a unique, previously unknown animal sound. Ecoacousticians have successfully enlisted the help of citizen scientists to gather data on bats (e.g., *Bat Detective*: www.batdetective.org) and birds (e.g., *eBird*: ebird.org) and to record soundscapes on a global scale (*Record the Earth*: www.recordtheearth.org). Italian sound artist David Monacchi's *Fragments of Extinction* project, initiated in 2001, records the world's undisturbed primary equatorial forests to highlight disappearing soundscapes and brings attention and urgency to the ongoing loss of a 'sonic heritage of millions of years of evolution' (Monacchi & Krause 2017). Ecological sound art is an effective medium for science dissemination, and immersive experiences of soundscapes can engage listeners on an emotional level. This acts as a powerful and accessible tool for inspiring public awareness about the value of ecoacoustics and ecosystem health in general (Monacchi & Krause 2017), and its efficacy in driving behavior changes is another interesting topic for scientific investigation.

A WAY FORWARD

With the increasing popularity of PAM and rapid flurry of analytical tools, it is now necessary to take advantage of obvious opportunities for acoustic data collection, to develop standards for data collection that allow cross-site comparisons and to create an open repository to store, visualize, and share recordings.

COLLECT MORE DATA.—Just as a meteorological station has become a standard and invaluable accessory at biological field stations, there should also be a permanent acoustic recorder. Anyone can put out a recorder, and researchers with long-term field programs are in a particularly good position to conduct passive acoustic monitoring for biodiversity. Long-term research sites typically have metadata related to vegetation composition and structure, faunal richness and abundance, and/or physical landscape variables that can be used together with acoustic data to create and validate population, community, and soundscape monitoring models. Detailed methods for collecting ecoacoustic data and a review of available hardware can be found in WWF's guide to acoustic monitoring (Browning *et al.* 2017); we encourage researchers to consult this report and take advantage of their field sites by beginning to compile invaluable long-term acoustic datasets that will contribute to creating a global database.

STANDARDIZED ACOUSTIC DATA COLLECTION.—To build a comprehensive PAM program, one needs to acquire the necessary hardware and software, develop a method for data collection, and determine a plan for storage of acoustic data files and associated metadata. While we understand that every PAM project will have specific requirements to address the research questions of interest, the best way to maximize the utility of any PAM effort is to follow a standard storage and metadata protocol. We strongly encourage researchers to use the data storage and metadata standards proposed by Roch *et al.* (2016). When acoustic data are organized and annotated in a uniform way, it allows other researchers (present-day or future) to utilize the data for additional questions.

A GLOBAL DATABASE.—With global data being increasingly publicly available in the ecological sciences (e.g., TRY, GBIF, GenBank, BOLD, eMammal), only a limited fraction meets the best practices standards defined by Joppa *et al.* (2016). Ideally, data should be freely available at high spatial resolution, up-to-date, user-friendly and assessed for accuracy, thereby increasing our ability to answer broad questions and improve its utility for conservation management. The Macaulay Library (<https://www.macaulaylibrary.org/>) and xeno-canto (<https://www.xeno-canto.org/>) are two large databases that house bioacoustic data, but only the latter allows full-soundscape recordings to be uploaded. Existing ecoacoustics databases include ARBIMON (<https://arbimon.sie-ve-analytics.com/home>), the Remote Environmental Assessment Laboratory (REAL, <http://lib.real.msu.edu/>), Ecosounds (<http://ecosounds.org>), and the Center for Global Soundscapes (<https://centerforglobalsoundscapes.org>), although only the first allows users to upload their own data. For marine acoustic data, there is support across US federal agencies to archive PAM recordings at the National Center for Environmental Information (NCEI); terrestrial and freshwater ecologists must follow suit. A free platform for soundscape storage to enable future temporal and spatial comparisons is absolutely necessary to advance tropical ecology and conservation.

CONCLUSION

We are convinced that PAM is a powerful tool that can be used to assess biodiversity over a range of spatial and temporal scales and can detect rare species, human impacts, and climatic shifts. Just as a plant or animal voucher specimen can provide information on diet, disease, and evolutionary relationships, so too can a sound recording provide information on species occurrence, density, distribution, phenology, inter- and intraspecific communication, and much more. The rapid proliferation of acoustic recorders and analysis algorithms makes this an exciting frontier in tropical ecology, yet we urge scientists to create standards in our approach to data collection, analysis, and archiving that will amplify the utility of recordings. What PAM can reveal will be invaluable in future decades as tropical ecosystems continue to change.

ACKNOWLEDGMENTS

We thank the participants and attendees of the ‘Quantitative acoustic ecology: Using sound in tropical biodiversity research and conservation’ symposium at the ATBC meeting in Merida, 2017, for insightful presentations and discussion of the need for documenting soundscapes in tropical ecosystems. We also thank two anonymous reviewers for providing feedback that improved the commentary. JLD was supported by the Center for Conservation and Sustainability at the Smithsonian Conservation Biology Institute; MCC acknowledges the support of the Brazilian Ministry of Education (CAPES); AKK was supported by the Max Planck Society; LDN was supported by the Brazilian National Council of Scientific and Technological Development (CNPq) 203230/2015-9 and Ecology Center USU; BCP has been supported by funds from Purdue University, the Office of the Executive Vice President for Research and Engagement, College of Agriculture Deans Office, NSF Coupled Natural and Human Systems Program, NSF Advancing Informal STEM Learning, and the USDA McIntire-Stennis Program; BLG has been supported by the Lynne Fellowship Program, Graduate School, Purdue University, and the Wright Forestry Fund, Department of Forestry and Natural Resources, Purdue University; SL was supported by the Australian Research Council DECRA DE130100565.

LITERATURE CITED

- AIDE, T. M., C. CORRADA-BRavo, M. CAMPOS-CERQUEIRA, C. MILAN, G. VEGA, AND R. ALVAREZ. 2013. Real-time bioacoustics monitoring and automated species identification. *PeerJ* 1: e103.
- AIDE, T. M., A. HERNÁNDEZ-SERNA, M. CAMPOS-CERQUEIRA, O. ACEVEDO-CHARRY, AND J. L. DEICHMANN. 2017. Species richness (of insects) drives the use of acoustic space in the tropics. *Remote Sens.* 9: 1096.
- ALVAREZ-BERRÍOS, N., M. CAMPOS-CERQUEIRA, A. HERNÁNDEZ-SERNA, C. J. AMANDA DELGADO, F. ROMÁN-DAÑOBEYTLA, AND T. M. AIDE. 2016. Impacts of small-scale gold mining on birds and anurans near the Tambopata Natural Reserve, Peru, assessed using Passive Acoustic Monitoring. *Trop. Conserv. Sci.* 9: 832–851.
- ASTARAS, C., J. M. LINDER, P. WREGE, R. D. ORUME, AND D. W. MACDONALD. 2017. Passive acoustic monitoring as a law enforcement tool for Afrotropical rainforests. *Front. Ecol. Environ.* 15: 233–234.
- BERTUCCI, F., E. PARMENTIER, G. LECHELLIER, A. D. HAWKINS, AND D. LECCHINI. 2016. Acoustic indices provide information on the status of coral reefs: An example from Moorea Island in the South Pacific. *Sci. Rep.* 6: 33326.
- BLUMSTEIN, D. T., D. J. MENNILL, P. CLEMINS, L. GIROD, K. YAO, G. PATRICELLI, J. L. DEPPE, A. H. KRKAUER, C. CLARK, K. A. CORTOPASSI, S. F. HANSER, B. MCCOWAN, A. M. ALI, AND A. N. G. KIRSCHER. 2011. Acoustic monitoring in terrestrial environments using microphone arrays: Applications, technological considerations and prospectus. *J. Appl. Ecol.* 48: 758–767.
- BORMPOUDAKIS, D., J. SUEUR, AND J. D. PANTIS. 2013. Spatial heterogeneity of ambient sound at the habitat type level: Ecological implications and applications. *Landscape Ecol.* 28: 495–506.
- BRIDGES, A. S., M. E. DORCAS, AND W. L. MONTGOMERY. 2000. Temporal variation in anuran calling behavior: Implications for surveys and monitoring programs. *Copeia* 2000: 587–592.

- BROWNING, E., R. GIBB, P. GLOVER-KAPFER, AND K. E. JONES. 2017. *Passive Acoustic Monitoring in Ecology and Conservation*. WWF-UK, Woking, UK.
- BURIVALOVA, Z., M. TOWSEY, T. BOUCHER, A. TRUSKINGER, C. APELIS, P. ROE, AND E. T. GAME. 2017. Using soundscapes to detect variable degrees of human influence on tropical forests in Papua New Guinea. *Conserv. Biol.* 1–11.
- BURTON, A. C., E. NEILSON, D. MOREIRA, A. LADLE, R. STEENWEG, J. T. FISHER, E. BAYNE, AND S. BOUTIN. 2015. Wildlife camera trapping: A review and recommendations for linking surveys to ecological processes. *J. Appl. Ecol.* 52: 675–685.
- BUTLER, J., J. A. STANLEY, AND M. J. BUTLER IV. 2016. Underwater soundscapes in near-shore tropical habitats and the effects of environmental degradation and habitat restoration. *J. Exp. Mar. Biol. Ecol.* 479: 89–96.
- BUXTON, R. T., E. BROWN, L. SHARMAN, C. M. GABRIELE, AND M. F. MCKENNA. 2016. Using bioacoustics to examine shifts in songbird phenology. *Ecol. Evol.* 6: 4697–4710.
- BUXTON, R., M. F. MCKENNA, M. CLAPP, E. MEYER, E. STABENAU, L. M. ANGELONI, K. CROOKS, AND G. WITTEMYER. 2018. Efficacy of extracting indices from large-scale acoustic recordings to monitor biodiversity. *Conserv. Biol.* <https://onlinelibrary.wiley.com/doi/abs/10.1111/cobi.13119>
- de CAMARGO, U. M., P. SOMERVUO, AND O. OVASKAINEN. 2017. PROTAX-Sound: A probabilistic framework for automated animal sound identification. *PLoS ONE* 12: e0184048.
- CAMPOS-CERQUEIRA, M., AND T. M. AIDE. 2016. Improving distribution data of threatened species by combining acoustic monitoring and occupancy modelling. *Methods Ecol. Evol.* 7: 1340–1348.
- CAMPOS-CERQUEIRA, M., AND T. M. AIDE. 2017. Lowland extirpation of anuran populations on a tropical mountain. *PeerJ* 5: e4059.
- CAMPOS-CERQUEIRA, M., W. J. ARENDT, J. M. WUNDERLE, AND T. M. AIDE. 2017. Have bird distributions shifted along an elevational gradient on a tropical mountain? *Ecol. Evol.* 7: 9914–9924.
- CORRADA BRAVO, C. J., R. ÁLVAREZ BERRÍOS, AND T. M. AIDE. 2017. Species-specific audio detection: A comparison of three template-based detection algorithms using random forests. *PeerJ Comp. Sci.* 3: e113.
- DARRAS, K., P. PÜTZ, K. REMBOLD, FAHRURROZI, AND T. TSCHARNTKE. 2016. Measuring sound detection spaces for acoustic animal sampling and monitoring. *Biol. Cons.* 201: 29–37.
- DEICHMANN, J. L., A. HERNÁNDEZ-SERNA, J. A. DELGADO C, M. CAMPOS-CERQUEIRA, AND T. M. AIDE. 2017. Soundscape analysis and acoustic monitoring document impacts of natural gas exploration on biodiversity in a tropical forest. *Ecol. Ind.* 74: 39–48.
- FULLER, S., A. C. AXEL, D. TUCKER, AND S. H. GAGE. 2015. Connecting soundscape to landscape: Which acoustic index best describes landscape configuration? *Ecol. Ind.* 58: 207–215.
- GASC, A., J. SUEUR, S. PAVOINE, R. PELLENS, AND P. GRANDCOLAS. 2013. Biodiversity sampling using a global acoustic approach: Contrasting sites with microendemism in New Caledonia. *PLoS ONE* 8: e65311.
- GEAY, T., P. BELLEUDY, C. GERVAISE, H. HABERSACK, J. AIGNER, A. KREISLER, H. SEITZ, AND J. B. LARONNE. 2017. Passive acoustic monitoring of bed load discharge in a large gravel bed river. *J. Geophys. Res., series F* 122: 528–545.
- GREGORY, T., F. C. RUEDA, J. DEICHMANN, J. KOLOWSKI, AND A. ALONSO. 2014. Arboreal camera trapping: Taking a proven method to new heights. *Methods Ecol. Evol.* 5: 443–451.
- HILL, A. P., P. PRINCE, E. PIÑA COVARRUBIAS, C. P. DONCASTER, J. L. SNADDON, AND A. ROGERS. 2018. AudioMoth: Evaluation of a smart open acoustic device for monitoring biodiversity and the environment. *Methods Ecol. Evol.* 9: 1199–1211.
- HOBBS, M. T., AND C. S. BREHME. 2017. An improved camera trap for amphibians, reptiles, small mammals, and large invertebrates. *PLoS ONE* 12: e0185026.
- JOPPA, L. N., B. O'CONNOR, P. VISCONTI, C. SMITH, J. GELDMANN, M. HOFFMANN, J. E. M. WATSON, S. H. M. BUTCHART, M. VIRAH-SAWMY, B. S. HALPERN, S. E. AHMED, A. BALMFORD, W. J. SUTHERLAND, M. HARFOOT, C. HILTON-TAYLOR, W. FODEN, E. D. MININ, S. PAGAD, P. GENOVESI, J. HUTTON, AND N. D. BURGESS. 2016. Filling in biodiversity threat gaps. *Science* 352: 416–418.
- KALAN, A. K., R. MUNDRY, O. J. J. WAGNER, S. HEINICKE, C. BOESCH, AND H. S. KÜHL. 2015. Towards the automated detection and occupancy estimation of primates using passive acoustic monitoring. *Ecol. Ind.* 54: 217–226.
- KRAUSE, B. 1987. Bio-acoustics: Habitat ambience and ecological balance. *Whole Earth Rev.* 57: 14–17.
- LINKE, S., E. DECKER, C. DESJONQUÈRES, AND T. GIFFORD. 1999. Temporal variation in underwater soundscapes: Implications for monitoring. *Freshw. Biol.* 42: 575–584.
- LINKE, S., T. GIFFORD, C. DESJONQUÈRES, D. TONOLLA, T. AUBIN, L. BARCLAY, C. KARACONSTANTIS, M. J. KENNARD, F. RYBAK, AND J. SUEUR. 2018. Freshwater ecoacoustics as a tool for continuous ecosystem monitoring. *Front. Ecol. Environ.* 16: 231–238.
- LLUSIA, D., R. MÁRQUEZ, J. F. BELTRÁN, M. BENÍTEZ, AND J. P. DO AMARAL. 2013. Calling behaviour under climate change: Geographical and seasonal variation of calling temperatures in ectotherms. *Glob. Change Biol.* 19: 2655–2674.
- MANKIN, R. W., D. W. HAGSTRUM, M. T. SMITH, A. L. RODA, AND M. T. K. KAIRO. 2011. Perspective and promise: A Century of insect acoustic detection and monitoring. *Am. Entomol.* 57: 30–44.
- MONACCHI, D., AND B. KRAUSE. 2017. Ecoacoustics and its expression through the voice of the arts: An essay. In A. Farina, and S. H. Gage (Eds.). *Eco Acoustics: The Ecological Role of Sounds*, pp. 297–311. Wiley, Hoboken, NJ.
- NARINS, P. M., AND S. W. F. MEENDERINK. 2014. Climate change and frog calls: Long-term correlations along a tropical altitudinal gradient. *Proc. R. Soc. B: Biol. Sci.* 281: 20140401.
- PIERETTI, N., A. FARINA, AND D. MORRI. 2011. A new methodology to infer the singing activity of an avian community: The Acoustic Complexity Index (ACI). *Ecol. Ind.* 11: 868–873.
- PIJANOWSKI, B. C., L. J. VILLANUEVA-RIVERA, S. L. DUMYAHN, A. FARINA, B. L. KRAUSE, B. M. NAPOLETANO, S. H. GAGE, AND N. PIERETTI. 2011. Soundscape Ecology: The science of sound in the landscape. *Bio-science* 61: 203–216.
- PLAISANCE, L., M. J. CALEY, R. E. BRAINARD, AND N. KNOWLTON. 2011. The diversity of coral reefs: What are we missing? *PLoS ONE* 6: e25026.
- RANKIN, L., AND A. C. AXEL. 2017. Biodiversity Assessment in Tropical Biomes Using Ecoacoustics: Linking Soundscape to Forest Structure in a Human-dominated Tropical Dry Forest in Southern Madagascar. *Ecoacoustics*, pp. 129–145. John Wiley & Sons, Ltd, Hoboken, NJ.
- RIBEIRO, J. W., L. S. M. SUGAI, AND M. CAMPOS-CERQUEIRA. 2017. Passive acoustic monitoring as a complementary strategy to assess biodiversity in the Brazilian Amazonia. *Biodivers. Conserv.* 26: 2999–3002.
- ROCH, M. A., H. BATCHELOR, S. BAUMANN-PICKERING, C. L. BERCHOK, D. CHOLEWIAK, E. FUJIOKA, E. C. GARLAND, S. HERBERT, J. A. HILDEBRAND, E. M. OLESON, S. VAN PARIJS, D. RISCH, A. ŠIROVIĆ, AND M. S. SOLDEVILLA. 2016. Management of acoustic metadata for bioacoustics. *Ecol. Inform.* 31: 122–136.
- RODRIGUEZ, A., A. GASC, S. PAVOINE, P. GRANDCOLAS, P. GAUCHER, AND J. SUEUR. 2014. Temporal and spatial variability of animal sound within a neotropical forest. *Ecol. Inform.* 21: 133–143.
- ROUNTREE, R. A., AND F. JUANES. 2017. Potential of passive acoustic recording for monitoring invasive species: Freshwater drum invasion of the Hudson River via the New York canal system. *Biol. Invasions* 19: 2075–2088.
- SCHAFFER, R. M. 1969. *The New Soundscape: A Handbook for the Modern Music Teacher*. Berandol Music Limited, Toronto, ON.
- SHONFIELD, J., AND E. M. BAYNE. 2017. Autonomous recording units in avian ecological research: current use and future applications. *Avian Conserv. Ecol.* 12: 1–13. <https://doi.org/10.5751/ACE-00974-120114>.
- STAATERMAN, E., M. B. OGBURN, A. H. ALTIERI, S. J. BRANDL, R. WHIPPO, J. SEEMANN, M. GOODISON, AND J. E. DUFFY. 2017. Bioacoustic measurements

- complement visual biodiversity surveys: Preliminary evidence from four shallow marine habitats. *Mar. Ecol. Prog. Ser.* 575: 207–215.
- SUEUR, J., S. PAVOINE, O. HAMERLYNCK, AND S. DUVAIL. 2008. Rapid acoustic survey for biodiversity appraisal. *PLoS ONE* 3: e4065.
- TAVOLGA, W. N. 2012. Listening backward: Early days of marine bioacoustics. In A. N. Popper, and A. D. Hawkins (Eds.). *The Effects of Noise on Aquatic Life*, p. 695. Springer-Verlag, New York, NY.
- THOMPSON, W. L. 2003. *Sampling Rare or Elusive Species: Concepts, Designs and Techniques for Estimating Population Parameters*. Island Press, Washington, DC.
- TONOLLA, D., V. ACUÑA, M. S. LORANG, K. HEUTSCHI, AND K. TOCKNER. 2010. A field-based investigation to examine underwater soundscapes of five common river habitats. *Hydrol. Process.* 24: 3146–3156.
- TOWSEY, M., L. ZHANG, M. COTTMAN-FIELDS, J. WIMMER, J. ZHANG, AND P. ROE. 2014. Visualization of long-duration acoustic recordings of the environment. *Procedia Comp. Sci.* 29: 703–712.
- VAN PARIJS, S. M., C. W. CLARK, R. S. SOUSA-LIMA, S. E. PARKS, S. RANKIN, D. RISCH, AND I. C. VAN OPZEELAND. 2009. Management and research applications of real-time and archival passive acoustic sensors over varying temporal and spatial scales. *Mar. Ecol. Prog. Ser.* 395: 21–36.
- VEGA, G., C. J. CORRADA-BRAVO, AND T. M. AIDE. 2016. Audio segmentation using Flattened Local Trimmed Range for ecological acoustic space analysis. *PeerJ Comp. Sci.* 2: e70.
- VILLANUEVA-RIVERA, L., B. PIJANOWSKI, J. DOUCETTE, AND B. PEKIN. 2011. A primer of acoustic analysis for landscape ecologists. *Landscape Ecol.* 26: 1233–1246.
- WREGE, P. H., E. D. ROWLAND, S. KEEN, AND Y. SHIU. 2017. Acoustic monitoring for conservation in tropical forests: Examples from forest elephants. *Methods Ecol. Evol.* 8: 1292–1301.
- XIE, J., M. TOWSEY, M. ZHU, J. ZHANG, AND P. ROE. 2017. An intelligent system for estimating frog community calling activity and species richness. *Ecol. Ind.* 82: 13–22.